

fire knowledge NETWORK bushfire CRC

PROGRAM B

→ Eucalypt decline in the absence of fire

Dugald Close and Neil Davidson
School of Plant Science, University of Tasmania

CRC UTAS

bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

→ Contributors to the project

1. Project Leader Neil Davidson (UTas)
2. Project Manager Dugald Close (UTas)
3. Forest Fire Management Group (national)
 - a) Department of Environment and Conservation (WA)
 - b) Forests NSW
 - c) Forestry Tasmania
 - d) Department of Sustainability and Environment (Vic)
 - e) Department of Environment and Heritage (SA)
 - f) Department of Arts, Environment and Heritage (Tas)
4. Tuart Health Research Group (Murdoch University, WA)

bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

→ Tasmanian Midlands: pre-European fire management
John Glover, Mills Plains 1834 C/O TMAG



bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire


→ Tasmanian Midlands: recent fire management (in the absence of grazing)
Photo: Fred Duncan, Forest Practices Authority



bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire


→ Coastal *E. gomphocephala* (Tuart) woodlands south of Perth, WA








bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

→ High Altitude *Eucalyptus delegatensis*: north-east and north-west Tasmania





bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

➤ **Soil nutrition**

- n = 6 sites, 10 trees per site, for each tree 4 sub-samples from top 5 cm of soil were bulked
- WA: Long unburnt had higher nitrate and total N
Tas: Long unburnt had higher ammonia and lower total P
- No differences in Zn, Mn, Fe or Mg

WA

Tas

bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

➤ **Tree foliar sampling**

bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

➤ **Tree foliar nutrition**

- n = 6 sites, 10 trees per site, 1 sub-sample per tree
- WA: Long unburnt had lower N and (not significant) P and lower Cu and Mn
Tas: Long unburnt had lower P
- No differences in other micronutrients

WA

Tas

bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

➤ **Effects of prescribed fire on soil nutrition (by ion exchange resin and seedling bioassay)**

bushfire CRC

PROGRAM B : Eucalypt decline in the absence of fire

➤ **Effects of prescribed fire on plant nutrient availability: WA**

- Plant nutrient availability by ion exchange resin
- n = 10 trees, 4 resin sacks (N-S-E-W under dripline) buried at 5 cm depth per tree
- N, P and all micronutrients (except Mn and Fe) higher in burn than no burn

bushfire CRC


PROGRAM B : Eucalypt decline in the absence of fire

➤ **Effects of prescribed fire on plant nutrient availability cont: WA**

- Plant nutrient availability by Tuart seedling bioassay
- n = 8 trees, 4 seedlings planted per tree
- Sampled early after 3 months due to kangaroo browsing
- P and Cu higher in burn than no burn seedling bioassays


Tree water use efficiency (by carbon isotope ratios): WA

1. n = 6 sites, 10 trees per site, 1 sub-sample per tree
2. Less negative ($p = 0.0379$) in long unburnt due to less water availability. Via competition with understorey?



Soil Microbial/Fungal Ecology

1. Average Well Colour Development (by Dr Yanfei Cai) as a measure of the functional diversity of the microbial communities (Garland and Mills 1991). Not a significant difference but trend toward higher functional diversity in state forest.
2. Tas: Sporocarp surveys of high altitude *E. delegatensis* sites (by Bryony Horton)
3. Observation indicates greater biodiversity in recently burnt compared to long unburnt sites. Molecular analysis being conducted in future.



Summary

1. Eucalypt decline - increased in the long absence of fire in some forest types (WA and Tas).
2. Decline is associated with the development of understorey, litter and decreased fungal diversity (WA and Tas).
3. Soil and Plant nutrition
 - WA
 - a) Increased N in long unburnt soil but lower N in long unburnt trees.
 - b) Micronutrients same in soils but Mn and Cu lower in long unburnt trees.
 - Tas
 - a) Increased N in long unburnt soil but same N in long unburnt trees.
 - b) Lower P in soil and far lower P in trees at long unburnt sites.

Summary cont.

4. Prescribed fire and plant nutrient availability (WA)
 - Resin exchange: Increased N, P and micronutrient (except Mn and Fe) availability.
 - Seedling bioassay: Increased P and Cu uptake on burn site, N and micronutrient uptake the same.
5. Trees have greater water use efficiency and are probably more water limited on sites with a long absence of fire (WA).

Conclusions - preliminary

1. Fire history is significantly associated with plant understorey and litter development, soil microbial diversity and activity, and tree health (decline) in some forest types in WA and Tas.
2. WA: Mn and Cu deficiency and limited water availability and; Tas: P deficiency are potential factors in eucalypt decline in the absence of fire.
3. Prescribed fire renders a large increase in plant nutrient availability (WA).

Future activity

1. Manipulative treatments of:
2. prescribed burning and;
3. understorey vegetation clearing
4. → for the restoration of tree health
5. = Conclusions from treatments (not correlations) on ecological processes that underpin eucalypt decline in the absence of fire



Acknowledgements

1. Funding and significant in-kind support (DEC and FT): Forest Fire Management Group
2. Fungal diversity work: Bryony Horton (UTas), Dr YanFei Cai (Murdoch Uni)
3. Industry reference group: Rick Sneeuwjagt (DEC), Drew Haswell (DEC), Tony Blanks (FT), Dr Tim Wardlaw (FT), Craig Brown (Forests NSW), Vic Jurskis (Forests NSW), Dr Paul Barber (Murdoch Uni)
4. Research plan advice: Professor Mark Adams
5. Technical assistance: Dr Perry Swanborough, Dr Robert Archibald, Peter Scott (Murdoch Uni), Chris Ware (UTas)
6. Statistical advice: Assoc. Prof. Ross Taplin (Murdoch Uni), Dr Greg Jordan (UTas)
7. Foliar sampling: Daan Lock (Arbor Oxygen), Dion Robertson (FT)